



THAWING OUT VIAL



1

Indicate the date and time when a vial of Pfizer-BioNTech COVID-19 Vaccine is removed from the freezer to thaw.


- Remove a thawed vial of the vaccine from the refrigerator and allow it to come to room temperature.
- If using a frozen vaccine, thaw for 30 minutes at room temperature.

Vials at room temperature must be diluted within 2 hours.

2

Prior to dilution
No more than
2 hours at room
temperature
(up to 25°C/77°F)

DILUTE BEFORE USE





3

- Before dilution, invert **gently** 10 times to mix.
- Do not shake.**

- Obtain sterile 0.9% Sodium Chloride Injection, USP.
- Cleanse the vial stopper with a single-use antiseptic swab.
- Add 1.8 mL of 0.9% Sodium Chloride Injection, USP into vaccine vial using a needle 21 gauge or narrower.
- Use a 3 mL syringe to draw up the diluent.

4




5

- Equalize vial pressure before removing the needle from the vial by withdrawing 1.8 mL air into the empty diluent syringe.

- Gently** invert the vial containing the vaccine to mix.
- Do not shake.**


6



7

- Record the date and time of dilution on the vaccine vial label.
- Store between 2°C to 25°C (35°F to 77°F).
- Discard any unused vaccine 6 hours after dilution.

PREPARATION OF INDIVIDUAL 0.3 ML DOSES OF PFIZER-BIONTECH COVID-19 VACCINE



8

- Using aseptic technique, cleanse the vial stopper with a single-use antiseptic swab, and withdraw 0.3 mL of vaccine.
- Using a 1 mL, 1 to 1 ½ inch, 25 gauge syringe to administer vaccine is preferred.
- Administer immediately, and no later than 6 hours after dilution.